



Tokyo Gakugei University Repository

東京学芸大学リポジトリ

<http://ir.u-gakugei.ac.jp/>

Title	被子植物の科の改良花式(abstract)
Author(s)	犀川, 政稔
Citation	東京学芸大学紀要. 自然科学系, 60: 69-75
Issue Date	2008-09-28
URL	http://hdl.handle.net/2309/91034
Publisher	東京学芸大学紀要出版委員会
Rights	

被子植物の科の改良花式

犀川 政稔*

環境科学分野

(2008年5月26日受理)

SAIKAWA, M.: Revised floral formulas of Angiosperm families. Bull. Tokyo Gakugei Univ. Natur. Sci., **60**: 69–75. (2008)

ISSN 1880-4330

Abstract

Floral formulas, as well as floral diagrams, are available for us to know morphological differences among Angiosperm taxa theoretically. In this study, the 114 floral formulas of families shown by Jones and Luchsinger (1987) were revised by removing and adding several American and Japanese families, respectively, resulting in a total of 122 floral formulas. Although floral formulas, such as those by Jones and Luchsinger have employed parentheses over and below the numerals and the abbreviations of calyx (CA), corolla (CO), androecium (A) and gynoecium (G) to show the union among these components of the flower, I have omitted the usage such parentheses in the floral formulas here mainly for computer users.

Key words: computer, floral formula, input, morphology

Department of Environmental Sciences, Tokyo Gakugei University, 4-1-1 Nukuikita-machi, Koganei-shi, Tokyo 184-8501, Japan

* 東京学芸大学環境科学分野 (184-8501 小金井市貫井北町 4-1-1)